arXiv:1511.01045v1 [math.GN] 2 Nov 2015

# Generating countable groups by discrete subsets

**Igor Protasov**

**Abstract.** Every countable topological group $G$ has a closed discrete subset $A$ such that $G = AA^{-1}$.

**MSC:** 22A05.

**Keyword:** topological group, generators.

## 1. Introduction

This note is to strengthen radically the following statement [1, Theorem 2.2]: every countable topological group can be generated by some closed discrete subset. All topologies under consideration are supposed to be Hausdorff.

**Theorem.** *Every countable topological group $G$ has a closed discrete subset $A$ such that $G = AA^{-1}$.*

*Proof.*We enumerate $G = \{g_n : n \in \omega\}$, $g_0 = e$, $e$ is the identity of $G$, and split the proof into two cases: $G$ is precompact (Case 1) and $G$ is not precompact (Case 2). We use two equivalent definitions of precompact groups: $G$ is a subgroup of some compact group and, for every neighborhood $U$ of $e$, there exists a finite subset $F$ of $G$ such that $G = FU$.

*Case 1.* Let $G$ be a dense subgroup of a compact group $H$ and let $\mu$ be the Haar measure on $H$. We choose a sequence $(r_n)_{n\in\omega}$ of positive real numbers such that $\Sigma_{i\in\omega} r_i < \frac{1}{6}$.

We put $x_0 = y_0 = e$, choose closed in $H$ neighborhood $U$ of $e$, such that $\mu(U) < r_0$, denote $U_0 = U$ and assume that, for some $n \in \omega$, we have chosen elements $x_0, \dots, x_n$ and $y_0, \dots, y_n$ of $G$, and closed in $H$ neighborhoods $U_0, \dots, U_n$ of $e$, such that , for each $k \leq n$ and $A_k = \{x_i, y_i : i \leq n\}$,

(1) $g_k \in A_k A_k^{-1}$;

(2) the subsets $\{x_i U_i : i \leq k\}$ are pairwise disjoint, the subsets $\{y_i U_i : i \leq k\}$ are pairwise disjoint and $x_i U_i \bigcap y_j U_j = \emptyset$ for each $i, j \in \{1, ..., n\}$;

(3) $\mu(U_i) < r_i$ for each $i \leq k$.

We suppose also that, for $n > 1$, there is a numeration $z_0, \dots, z_{m(n)}$ of $A_n A_n^{-1} \setminus A_n$ and closed in $H$ neighborhoods $V_0, \dots, V_{m(n)}$ of $e$ such that , for each $k \in \{1, \dots, n\}$, $A_k A_k^{-1} \setminus \{A_k \bigcup A_{k-1} A_{k-1}^{-1}\} = \{z_{m(k-1)}, ..., z_{m(k)}\}$, and

(4) $z_i V_i \bigcap A_k = \emptyset$, $i \in \{0, ..., m(k)\}$;

(5) $\mu(v_i) < r_i$, $i \in \{0, ..., m(k)\}$.

To make the inductive step from $n$ to $n+1$, we take the first element $g \in \{g_i : i \in \omega\} \setminus A_n A_n^{-1}$ and denote

$$B = \{x_i U_i : i \leq n\} \bigcup \{y_i U_i : i \leq n\} \bigcup \{z_i V_i : i \leq m(n)\}, \quad C = H \setminus B.$$

By (3), (5) and the choice of $(r_n)_{n\in\omega}$, we have $\mu(C) > \frac{1}{2}$. Hence, $gC \bigcap C \neq \emptyset$. Since $C$ is open in $H$, there are $x, y \in G \bigcap C$ such that $x = gy$. We put $x_{n+1} = x$, $y_{n+1} = y$, $A_{n+1} = A_n \bigcup \{x_{n+1}, y_{n+1}\}$ and note that $g \in A_{n+1}A_{n+1}^{-1}$. Since $x_{n+1}, y_{n+1} \in G \setminus B$, there is a closed in $H$ neighborhood $U_{n+1}$ of $e$ such that $x_{n+1}U_{n+1} \bigcap B = \emptyset$, $y_{n+1}U_{n+1} \bigcap B = \emptyset$, $x_{n+1}U_{n+1} \bigcap y_{n+1}U_{n+1} = \emptyset$ and $\mu(U_{n+1}) < r_{n+1}$. Hence, (1), (2) (3) are satisfied for $k = n + 1$.

We put $z_{m(n)+1} = y$ and enumerate $z_{m(n)+1}, ..., z_{m(n+1)}$ the set $A_{n+1}A_{n+1}^{-1} \setminus (A_{n+1} \bigcup \{z_0, ..., z_{m(n)}\})$. Then we choose closed in $H$ neighborhoods $V_{m(n)+1}, ..., V_{m(n+1)}$ of $e$ such that $z_iV_i \bigcap A_{n+1} = \emptyset$, $\mu(V_i) < r_i$ for each $i \in \{m(n)+1, ..., m(n)\}$. Thus, (4), (5) are satisfied for $k = n+1$.

After $\omega$ steps we get the desired $A = \{x_n, y_n : n \in \omega\} : G = AA^{-1}$ by (1), $A$ is discrete by (2), $A$ is closed by (4).

*Case 2.* We take a neighborhood $U$ of $e$ such that $G \neq FU$ for every finite subset $F$ of $G$, and pick a neighborhood $V$ of $e$ such that $VV^{-1} \subset U$. Then we choose inductively a sequence $(x_n)_{n\in\omega}$ in $G$ such that $\{x_n, g_nx_n\}V \bigcap \{x_i, g_ix_i\}V = \emptyset$ for each $i < n$. The set $A = \{x_n, g_nx_n : n \in \omega\}$ is closed, discrete and $G = AA^{-1}$.

**Question.** *Can every countable topological group $G$ be factorized into two (close) discrete subsets $A$ and $B$: $G = AB$ and the subsets $\{aB : a \in A\}$ are pairwise disjoint?*

**Remark.** A subset $A$ of a group $G$ is called *thin* if $gA \bigcap A$ is finite for each $g \in G \setminus \{e\}$. Can every countable topological group be generated by some thin closed discrete subset [2, Question 5]? In both cases of the proof, we have infinitely many possibilities to prolong $A_n$ to $A_{n+1}$, so the set $A$ in Theorem can be chosen thin.

## References


[1] W. Comfort, S. Morris, D. Robbie, S. Svetlichny, M. Tkachenko, *Suitable sets for topological groups*, Topology Appl. **86** (1998) 25-46.

[2] I. Protasov, *Thin subset of topological groups*, Topology Appl. **160** (2013), 1083-1087.



Department of Cybernetics, Kiev University.
Prospect Glushkova 2, corp. 6,
03680 Kyiv, Ukraine
e-mail: I.V. Protasov@gmil.com